

## Original article

### Pathological locations of uterine pregnancies

#### Patologické lokalizace gravidity v děloze

*Sergiu Leahomschi, Zdeněk Žížka, Zdeňka Fartáková*  
Fetal Medicine Center, Department of Obstetrics and Gynecology, 1st Medical Faculty,  
Charles University, Prague

Correspondence: MUDr. Sergiu Leahomschi, Gynekologicko-porodnická klinika 1. LF UK a VFN, Apolínářská 18, 128 08 Praha 2, tel.: +420 608 321 557, e-mail: sergiu.leahomschi@gmail.com

Published: 10. 10. 2014      Received: 3. 10. 2014      Accepted: 7. 10. 2014  
Actual Gyn 2014, 6, 81      ISSN 1803-9588      © 2014, Aprofema s.r.o.  
Free fulltext article at [www.actualgyn.com](http://www.actualgyn.com)



Cite as: Leahomschi S, Zizka Z, Fartakova Z. Pathological locations of uterine pregnancies. Actual Gyn. 2014;6:81

Uterine locations of ectopic pregnancies (interstitial, cervical and in the scar after Cesarean section) represent a minor part of all ectopic pregnancies, however their diagnosis is complicated and therefore their mortality is much bigger when compared to extrauterine ectopic pregnancies. On time diagnosis and adequate management of these cases are important goals for ultrasonographers involved in gynecological field.

In this work we pay attention to correct nomenclature, especially in the case of so-called "angular pregnancy" – we believe this term is obsolete. Interstitial location of

ectopic pregnancy should be differentiated from asymmetrically located normal intrauterine pregnancy. We propose a list of criteria for ultrasound diagnostics of pathologically located uterine pregnancies. Conservative therapy with methotrexate, its indications, contraindications and protocols are discussed. Surgical procedures were shortly described using a video presentation. Finally, we present a list of selected uterine ectopic pregnancies managed in our department (n = 12) describing the management for each case.